Range and Livestock Grazing

Specialist Report

East Fork Boulder Creek Native Trout Restoration Project USDA-Forest Service-Dixie National Forest

Date: June 30, 2011

Submitted by: /s/ Ron Mortensen and /s/ Georgina Lampman

Ron Mortensen, Rangeland Management Specialist Dixie National Forest Escalante Ranger District

and

Georgina Lampman Regional Planner Intermountain Regional Office



The following is an analysis of effects of the proposed East Fork Boulder Creek Native Trout Restoration Project on range and livestock grazing. The treatment area is displayed in Figure A. The alternatives that are analyzed in detail, including actions that are not part of the Forest Service decision but connected to the project, are described in Appendix 1.

Description of Affected Environment and Analysis Method

The project area includes the Boulder Allotment, administered by the Forest Service. The treatment streams are used as a water source for the livestock on the allotment. Riparian and upland vegetation are used as forage by livestock.

The overall area of analysis for direct and indirect effects on livestock grazing is the Between the Creeks Unit (pasture) of the Boulder Allotment (see Figure B). This pasture is expected to be grazed from approximately July 1 thru August 14 during years 2011 and 2013 and August 15 thru October 1 during years 2012 and 2014. The alternating periods of use is expected to continue.

Direct effects:

The direct effects from the proposed project are as follows:

- The potential direct effect on livestock grazing is the availability of water, either because of potential changes in water quality or water quantity.
 - o The effect will be measured by the miles of stream available for water.
 - The area of analysis is the Between the Creeks pasture, Boulder Allotment.
- The potential direct effect on livestock grazing is the availability of vegetation for forage, either because of potential damage to vegetation during or from treatment.
 - The effect will be measured by the amount of vegetation affected.
 - The area of analysis is the Between the Creeks pasture, Boulder Allotment.
- The potential direct effect to range vegetative communities, mountain meadows and parks with sage-grass or grass-forb vegetation, would be potential damage to vegetation during or from treatment.
 - The effect will be measured by the amount of vegetation affected.
 - The area of analysis is the Between the Creeks pasture, Boulder Allotment.

Indirect effects:

No indirect effects are anticipated.

Cumulative effects:

- The potential cumulative effects on livestock grazing would be past, present, and reasonably foreseeable actions that affect the availability of water and vegetation.
 - The effect will be measured as for the direct effects.
 - o The cumulative effects area of analysis is the Boulder Allotment.

Direct and Indirect Effects

No Action

Under the No Action alternative, there would be no treatment; therefore, there would be no change in and thus no effect to water quality or quantity, vegetation for forage, or range vegetative communities.

Proposed Action

The Proposed Action would result in rotenone application to 7.4 miles of stream within the Between the Creeks pasture of the Boulder Allotment. As described in the Floodplains, Wetlands, and Water Quality specialist report, there would be no effect to water quality relating to irrigation water and stock water; therefore, there would be no effect to water availability for livestock. As described in the Forest Service risk assessment of rotenone, "there is no basis for asserting that adverse effects are plausible in large or small mammals when rotenone is applied at the highest application rate considered in this risk assessment, 200 ppb" (Durkin 2008). Rotenone application for the Proposed Action is expected to be less than the maximum and would decrease as the chemical degrades, often in less than 24 hours, as described in the specialist report on Water Quality, Floodplains, and Wetlands. Livestock would also be exposed to potassium permanganate in the 0.25 to 0.5 miles below the neutralization site at decreasing concentration until the potassium permanganate oxidizes the rotenone. At the rate it would be applied, 2-6 mg/l, potassium permanganate would not affect water quality for livestock and would be decreasing in concentration through the 0.25 to 0.5 mile long neutralization zone until it oxidizes the rotenone. No adverse acute effects to livestock are expected from the chemical. Because potential exposure of livestock to the chemicals would occur only if treatment were to occur prior to October 1, either 1 or 2 years out of the potential 3 years of treatment, and for only 24 hours, no chronic exposure or effects would occur. Other water is also available for livestock in the area.

The Proposed Action would involve access by personnel setting up application and neutralization sites. Because access to application sites would be by foot, and application would be done with drip stations, the effects on riparian and upland vegetation would be minimal. Access to neutralization sites may be by motorized vehicle and potentially affect vegetation if cross-country travel is needed, but this would be minimal and occur at only the lower end of the treatment areas. Overlap in period of use by livestock and disturbance to forage would be for short duration, and vegetation is available for livestock elsewhere in the pasture.

The effects to range vegetative communities under the Proposed Action are as described for effects to livestock forage availability. It would be short duration and minimal. Design criteria require application procedures to be in accordance with policy, which includes the Forest Noxious Weed strategy. and UDWR would be expected to follow the Utah Noxious Weed Act requirements regarding treatment of machinery in Utah State Code R68-9-4; therefore, there would also be no expected increase in noxious weeds to affect range vegetative communities from the Proposed Action.

Non-chemical Treatment

As described in the specialist report on Floodplains, Wetlands, and Water Quality, the Non-chemical Treatment alternative would result in short term fine sediment in the 7.4 miles of stream within the Between the Creeks pasture of the Boulder Allotment. This would not affect the quality of water for livestock. Water quality and quantity for livestock would not be affected.

The Non-chemical Treatment alternative would involve multiple passes of personnel conducting electrofishing that includes hiking along edges and sweeping through streambank and aquatic vegetation with electrofishing equipment. Riparian vegetation could be disturbed in the process along the 7.4 miles of stream in the pasture. Effects would be short-term, because the vegetation should recover by the following growing season. Overlap in period of use by livestock and disturbance to vegetation would be for short duration in any one location as crews move through. Overall presence would be approximately 40 days per year, based on an assumption that only two removal efforts would occur during the period the pasture would be in use. This may occur for up to 10 years. Other forage would be available for livestock elsewhere in the pasture.

The effects to range vegetative communities under the Non-chemical Treatment alternative are as described for effects to livestock forage availability. It would be short duration and minimal. UDWR would also be expected to follow the Utah Noxious Weed Act requirements regarding treatment of machinery in Utah State Code R68-9-4; therefore, there would be no expected increase in noxious weeds from the Non-chemical Treatment alternative.

Cumulative Effects

Under all alternatives, the effects to water availability and forage availability would be either non-existent, too small to be measured, or of too short duration relative to the pasture use period to have measurable effect; therefore, there would be no cumulative effects to water availability for livestock.

Under all alternatives, the effects to vegetation would be either non-existent, too small to be measured, or of too short duration relative to the pasture use period to have measurable effect; therefore, there would be no cumulative effects to vegetation for livestock forage.

Summary

There would be no direct, indirect, or cumulative effects to range resources, livestock, and noxious weeds from the No Action alternative. There would be non-existent, too small to measure, or of too short duration relative to the pasture use period to have measurable direct effects to range resources, livestock, and noxious weeds under the Proposed Action and non-chemical alternative. There would be no indirect or cumulative effects to range resources or livestock.

Compliance with Other Laws and Regulations

All alternatives are compliant with laws and regulations affecting livestock grazing and management.

Forest-plan Consistency Determination

All alternatives are consistent with Forest Plan direction.

Use and/or consideration of Best Available Science

This analysis considered the water quality analysis for the project, which in turn considered the best available science on effects of rotenone and potassium permanganate on non-fish organisms. Analysis of effects to the vegetation considered knowledge based on effects to riparian vegetation from recreation and other minimal disturbance activities on this and other allotments of the District.

Literature Cited

Durkin, P.R. 2008. Rotenone: Human health and Ecological Risk Assessment, Final Report. USDA Forest Service Contract AG-3187-C-06-0010. 217 pp.

Figure A. East Fork Boulder Creek Native Trout Restoration project area





Figure B. Livestock grazing allotments in vicinity of proposed East Fork Boulder Creek Native Trout Restoration Project



Appendix 1. Project Area and Alternatives Analyzed in Detail

The following describes and compares the Forest Service alternatives analyzed. It includes a description of the UDWR's proposed project and considers UDWR's treatment alternative in detail. This section also presents the alternatives and the UDWR activities that would be authorized or connected actions to the alternatives in comparative form.

Project Area

The proposed East Fork Boulder Creek Native Trout Restoration Project (project) is located approximately 7 miles northwest of Boulder, Utah (see Figure 1). The total treatment area is as follows:

- approximately 7.8 miles (12.6 km) of East Fork Boulder Creek from the natural barrier (below headwater meadow) on East Fork Boulder Creek to its confluence with West Fork Boulder Creek;
- approximately 0.2 miles (0.4 km) of lower West Fork Boulder Creek, from a previously constructed barrier to its confluence with East Fork Boulder Creek;
- approximately 0.5 miles (0.8 km) of Boulder Creek from the confluence of East Fork Boulder Creek and West Fork Boulder Creek downstream to a previously constructed fish barrier;
- all seeps and springs flowing into those sections of streams proposed for fish removal; and
- the Garkane Energy water transfer pipeline between the West Fork Reservoir and King's Pasture Reservoir; King's Pasture (East Fork) Reservoir; a pond on private property in King's Pasture, and the Garkane Energy penstock, between King's Pasture Reservoir and the Garkane Energy Boulder Creek Hydroelectric Power Plant (main power plant).

The treatment stream reaches flow through portions of Sections 27, 28, 33, and 34 of T31S, R4E, and Sections 3, 10, 15, 21, 22, and 28 of T32S, R4E, Salt Lake Baseline Meridian. Treatment would include connecting waters, including relatively large inflows or tributaries with permanent fish habitat and smaller springs and seeps that are capable of at least temporarily holding small fish. Known tributaries and inflows vary in length from 10 meters to over 750 meters.

The reaches on NFS-lands are all on the Escalante Ranger District of the Forest in Garfield County, Utah. The inflow of the water transfer pipeline is at the West Fork Reservoir in Section 8, T32S, R4E, and the outflow is at King's Pasture Reservoir in Section 10 of T32S, R4E. The inflow of the penstock is at King's Pasture Reservoir, and the outflow is at the main power plant in Section 35 of T32S, R4E.

No Action- No Further Treatment Scenario

Under the No Action alternative, the Forest would not approve the pesticide use permit to UDWR, would not authorize UDWR to use motorized vehicles off of designated routes for the application of rotenone to waters of the treatment area on NFS lands, and would not approve a special use authorization for UDWR to bury removed fish.

The No Action alternative would not preclude UDWR from implementing actions on NFS lands that would meet the purpose and need for UDWR's project but do not require Forest Service authorization. This includes UDWR activities described under the Non-chemical Treatment alternative (Section 2.1.3) except for the use motorized vehicles off of designated routes or burial of removed fish on NFS lands. The No Action alternative would also not preclude UDWR from implementing actions on non-NFS lands that are related to the purpose and need for UDWR's project but not under Forest Service jurisdiction or authorization.

One possible option for UDWR is to take no further action to meet the purpose and need of the proposed project. This possible option is identified in this analysis as the "No Action - No Further Treatment Scenario" and is the basis for the effects analysis for the No Action alternative to provide the base line for comparison of expected future conditions if neither the Proposed Action nor Non-chemical Treatment alternative were implemented by the Forest and UDWR were to take no further action to meet the purpose and need.

Proposed Action

The Proposed Action is to approve the pesticide use permit that the Forest Service requires the UDWR to have to apply the fish toxicant rotenone to waters that flow on NFS lands and to authorize motorized vehicle use off of designated routes. The pesticide use permit would authorize the UDWR to implement a maximum of three treatments on NFS land, one treatment per year for three consecutive years. Waters on NFS land that would be treated by UDWR under the Forest Service pesticide use permit are as follows:

- approximately 7.8 miles (12.6 km) of East Fork Boulder Creek from the natural barrier (below headwater meadow) on East Fork Boulder Creek to its confluence with West Fork Boulder Creek;
- approximately 0.2 miles (0.4 km) of lower West Fork Boulder Creek, from a previously constructed barrier to its confluence with East Fork Boulder Creek;
- approximately 0.5 miles (0.8 km) of Boulder Creek from the confluence of East Fork Boulder Creek and West Fork Boulder Creek downstream to a previously constructed fish barrier; and
- all seeps and springs flowing into those sections of the stream reaches specified in the permit.

The UDWR activities that would be authorized by the Forest under the Proposed Action would completely eradicate non-native trout from East Fork Boulder Creek, a short segment of Boulder Creek, and a very short segment of West Fork Boulder Creek. All fish would be temporarily eliminated by UDWR from target waters. Use of motorized vehicles by UDWR off of designated routes may be needed to facilitate placement of equipment, especially neutralization equipment, in effective locations.

Several actions that are not part of the Forest Service decision are connected to the UDWR project, as follows. UDWR is proposing chemical treatment of connected waters on private property to meet the purpose of the UDWR project. Following fish removal, UDWR would introduce the CRCT into the treated stream segments to establish self-sustaining populations. Sterile hybrids of species of non-native trout may also be stocked by UDWR at some locations following the treatments to provide sport fishing

opportunities while native trout become established. The following describes the UDWR project in detail, including identification of those actions that do not require Forest Service authorization.

Chemicals. Liquid emulsifiable rotenone (Liquid Rotenone, 5% Active Ingredient, EPA Registration No. 432-172) would be used by UDWR to treat target waters. Rotenone was selected as the chemical to use because of its effectiveness in controlling fish populations and its lack of long-term effects on the environment (Sousa et al 1987). When used at the concentrations planned for the UDWR project, rotenone is a naturally occurring fish toxicant that is toxic to only fish, some aquatic invertebrates, and some juvenile amphibians. EPA found it to be not toxic to humans, other mammals, and birds at the concentrations used to remove fish (EPA 2007). It has been widely used in the United States since the 1950's. UDWR has used rotenone successfully in many similar projects and has refined application techniques to minimize adverse side effects to the environment (Hepworth et al. 2001a, Hepworth et al. 2001b, Hepworth et al. 2001c, Ottenbacher and Hepworth 2001, Chamberlain and Hepworth 2002a, Chamberlain and Hepworth 2002b, Chamberlain and Hepworth 2002c, Fridell et al. 2004, Fridell et al. 2005, Fridell and Rehm 2006).

Potassium permanganate would be used by UDWR to neutralize the rotenone at suitable locations to prevent the movement of rotenone into non-target waters. Potassium permanganate was selected, because it is a strong oxidizer that breaks down into potassium, manganese, and water. All are common in nature and have no deleterious environmental effects at the concentrations that would be used for the UDWR project activities, including those that would be authorized by the Forest under the Proposed Action (Finlayson et al. 2000). Potassium permanganate is used as an oxidizing agent in treatment plants to purify drinking water (EPA 1999). Although the oxidation process is not immediate, neutralization should occur within an estimated 0.25 to 0.5 miles of the neutralization site.

A more detailed description of the chemicals that would be used for the UDWR project activities, including those that would be authorized by the Forest under the Proposed Action, can be found in specialist report on Chemicals and Application of the Proposed Action.

Application. Liquid rotenone would be applied by UDWR at a rate of 0.5 to 2.0 ppm. In the pond and reservoir, liquid rotenone would be dispersed from personnel on small water-craft using pressurized backpack spray units. For flowing waters, seeps, and springs, liquid rotenone would be applied using a combination of 30 gallon and 5 gallon dispensers with constant flow drip-heads at approximately 50 to 60 stations throughout the UDWR project area over a 3 to 24 hour period (Finlayson et. al 2000, Ottenbacher et al. 2009). One 30 gallon drip station would be used by UDWR at each at the following:

- lower end of the headwater meadow at the upstream end of the UDWR project area,
- approximately halfway between the headwater meadow and King's Pasture Reservoir,
- immediately below King's Pasture Reservoir, and
- at the intake for the water flow pipeline between the West Fork Reservoir and King's Pasture Reservoir.

Five-gallon drip stations would be located by UDWR at approximately 1 mile intervals, beginning one mile below King's Pasture Reservoir and ending 1 mile upstream from the fish barriers on the main stem of East Fork Boulder Creek, and at all major springs and seeps within the UDWR project area. The

interval placement of drip stations on the main stem of East Fork Boulder Creek would be to facilitate efficient travel time of chemicals. Depending on flow volume, a single 30 gallon or 5 gallon drip would be placed by UDWR on the lower fish barrier on West Fork Boulder Creek. Pressurized backpack sprayers would be used by UDWR to apply a diluted solution of the chemical to springs and backwater areas containing fish that were not effectively treated by boat or drip station.

Rotenone would be neutralized by UDWR with potassium permanganate downstream from target waters. Three sites are planned: where the penstock water is released at the upper power plant, where water is released at the main power plant, and at the fish barrier at the lower end of the treatment area. Each site would have a main neutralization station and at least one contingency neutralization station to ensure effectiveness. The neutralization stations would prevent rotenone from escaping the target area, except for the estimated 0.25 to 0.5 miles downstream in which the neutralization or natural degradation of rotenone would be occurring.

Post-treatment activity. Following confirmation of complete non-native trout removal, UDWR would reintroduce CRCT into project stream reaches from "core" CRCT populations or from fish produced by UDWR CRCT brood stocks. Sterile hybrids of species of non-native trout may also be stocked by UDWR at some locations following the treatments to provide sport fishing opportunities while native trout become established. All UDWR transfers or stocking of fish would comply with Utah Department of Agriculture and Food rules and UDWR policies.

Design Criteria. The following design criteria would be implemented and included in the Forest Service authorizations:

- 1. Stream sections will be treated in the fall to minimize impacts on non-target wildlife species (amphibians, insectivorous birds and bats). The fall treatment period will also minimize the impacts on sport fishing recreation.
- Each treatment will be preceded by internal and external notifications and media releases to notify
 the public of treatment sites and dates and will include the following: notification of the Boulder
 Town Council, notification of private landowners in the treatment area, and news releases in local
 papers.
- 3. The treatment area will be placarded to prohibit public access during treatment and for at least 3 days following treatment.
- 4. Application of the chemical will be conducted by licensed pesticide applicators in accordance with all applicable regulations and policies.
- 5. Access by motorized vehicles will be on National Forest System roads designated for motorized vehicle use to the extent possible. Any use of motorized vehicles off of designated routes will be minimal and will require written Forest Service approval.
- 6. Neutralization sites will be placed to maximize their effectiveness at preventing downstream escapement of rotenone.
- 7. Treated waters will remain open to fishing.

- 8. Transport to the site and storage of chemicals on the site will comply with FSH 2109.14.40 (Pesticide-Use Management and Coordination Handbook, Chapter 40 Storage, Transportation, and Disposal).
- 9. Sentinel fish ("in situ bioassay") will be used for pesticide residues monitoring to determine the presence or absence of unacceptable environmental effects.
- 10. Treatments will be discontinued if the objective of complete removal of non-native trout from the project area has been met.

Actions connected to but not included in the decision. The following parts of the UDWR project, as described above, are not subject to Forest Service permit requirements, and therefore are not included in the Forest Service decision. Selection of the Proposed Action is for issuance of the pesticide use permit for the application of rotenone on NFS lands only. The following, however, are considered connected actions and thus included in the environmental analysis:

1. The proposed UDWR treatment area includes private property, including property owned by Garkane Energy; thus, this area is not under Forest Service jurisdiction. This includes approximately 1.4 miles of East Fork Boulder Creek, Kings Pasture Reservoir, and the pond in Kings Pasture. To meet the purpose and need of the UDWR project, these areas as well as the water in the transmission pipeline and penstock must be treated by UDWR. Forest Service approval of the pesticide use permit for UDWR to apply rotenone to waters on NFS land is not approval of UDWR activities on non-NFS lands; however, the Forest Service would not approve the pesticide use permit unless UDWR is able to complete its project by treating waters off of NFS land.

The expectation is that the entire UDWR project treatment area would receive chemical treatment as described below, although the UDWR may decide to use another method or methods to achieve the treatment objective. FERC license order Section 4(e), item 16, condition 4, requires Garkane Energy to use its reasonable efforts to cooperate in the work of UDWR and other agencies to remove non-native fish and re-establish CRCT in the above stream sections. This cooperation has already been demonstrated through construction of the fish barriers and through the first chemical treatment of Kings Pasture Reservoir in 2009.

2. Stocking of fish is under the jurisdiction of UDWR; thus, the CRCT stocking is not under Forest Service jurisdiction. To meet the purpose and need of the UDWR project, the stream would need to be stocked by UDWR with CRCT from core populations or UDWR brood stock post-treatment.

The expectation is that the post-treatment recolonization/stocking of CRCT would occur as described. The purpose and need for the UDWR project, including stocking with CRCT, is to implement conservation actions under the CRCT Conservation Agreement and Strategy, to which UDWR is a signatory. In addition, the Forest Service conditions regarding the non-native fish eradication and fish restocking were included in a 2006 settlement agreement relating to the FERC license conditions and signed by Garkane Energy, Forest Service, and UDWR.

3. Fishing regulations, including whether or not treated waters would remain open to fishing, is under the jurisdiction of UDWR.

The expectation is that UDWR would manage the fishing regulations to meet the conservation actions under the CRCT Conservation Agreement and Strategy. UDWR recognizes the importance of the area to recreation users. Because of this, UDWR may also stock sterile hybrids of species of non-native trout at some locations following the treatments while native trout become established.

Non-chemical Treatment Alternative

Under the Non-chemical Treatment alternative, the Forest Service would authorize UDWR to use motorized vehicles off of designated routes and approve a special use authorization for UDWR to bury fish that are removed as necessary to implement a non-chemical treatment to remove non-native trout from waters on NFS land.

The non-chemical treatment methods would not involve the use of rotenone or other pesticides on NFS lands and, therefore, would not require Forest Service approval. The effects of the non-chemical treatment are being analyzed, because this option may be exercised by UDWR in the event that the Forest Service were to choose not to authorize pesticide use, and the approach would be a connected action to the authorization of the use of motorized vehicles off of designated routes and approval of a special use authorization for burial of removed fish. The other connected actions that would also not require new Forest Service action are described below. UDWR's non-chemical treatment and other connected actions may or may not occur under the No Action alternative if the UDWR were to use motorized vehicles only on designated routes. These UDWR actions also may or may not occur under the Proposed Action.

Under the Non-chemical Treatment alternative, UDWR would use electrofishing to remove non-native trout from the treatment waters on NFS lands. Except for possible motorized vehicle use off of designated routes and burial of removed fish, this alternative would not require Forest Service authorization.

Treatment area. The treatment area would remain the same as described in the Proposed Action.

Methodology and Equipment. Electrofishing would be used by UDWR to remove non-native trout from the treatment area on NFS lands. Electrofishing introduces an electric current into the water and is commonly used as a fish removal method. The electricity causes an involuntary muscle contraction in the fish, attracting them toward the source of the electricity (electrode). Workers with long-handled nets then collect the stunned fish. Voltage, amperage, pulse frequency, and waveform are manipulated to maximize effectiveness, which can be influenced by water flow and velocity, temperature, clarity, conductivity (dissolved mineral content), and substrate. Other factors influencing effectiveness include the fish size, species and behavior, presence of aquatic vegetation, time of year, and time of day. It is most effective in shallow water and is, therefore, most commonly used in rivers and streams and occasionally in the shallow water zones of lakes.

Electrofishing removal would be accomplished by UDWR using multiple Smith-Root LR24 backpack electrofishing units or their equivalent from another manufacturer. Block nets of sufficient width would be set up to prevent fish emigration during removal activities. Dip nets, buckets, and live wells would also be necessary for capture and removal of brook trout (*Salvenlinus fontinalis*) and capture and safe holding of CRCT.

Removal activities. Mechanical removal of non-native trout species using backpack electrofishing has been attempted in several other projects (Moore et al. 1986, Meronek et al. 1996, Thompson and Rahel 1996, Buktenica et al. 2000, Kulp and Moore 2000, Shepard et al. 2002, Peterson et al. 2004, Moore et al. 2005, Meyer et al. 2006, Earle et al. 2007). The results of these prior mechanical removal projects indicate: 1) achieving complete mechanical removal of trout in streams with the width, complexity, and number of small, heavily vegetated springs/tributaries found in East Fork Boulder Creek would be difficult; 2) success would be enhanced by implementing multiple-pass depletion removal efforts 3 to 4 times within the same year, and 3) success would be enhanced by treatment over multiple years (minimum of 2). For this UDWR project, the multi-year removal effort would involve a minimum of 5 to 6 people conducting multiple-pass removal efforts for the majority of summer and early autumn (late June to September) over a period of several years. While such removal efforts would undoubtedly cause major reductions in brook trout density and biomass, they may or may not result in complete eradication. UDWR would begin CRCT reintroduction efforts only when no brook trout are found within the project area.

The electrofishing removal by UDWR would follow the population monitoring methods used by Utah State University's Institute for Natural Systems Engineering, Utah Water Research Lab (INSE) during their Garkane-funded fish population monitoring on the Boulder Creek system (Hardy et al. 2009a, Hardy et al. 2009b). Personnel would electrofish approximately 100-meter reaches in 8.5 miles of the mainstem of East Fork Boulder Creek, West Fork Boulder Creek, and Boulder Creek along with all spring inflows and tributary streams. A block net would be placed across the upstream and downstream end of each reach to increase capture efficiency by preventing emigration. Up to 4 passes, or until no fish were collected, would be completed through each reach. Each pass would involve all personnel walking in the stream channel and on the banks while applying constant electric current to the water from at least two backpack electrofishers. All organisms within the stream would be subjected to the electric field. All non-native brook trout would be removed from the system, killed and buried. Any CRCT collected would be held in buckets/live wells and returned to the stream after completion of the 4 pass removal.

Effort. One crew would consist of at least 2 personnel using backpack electrofishers, 2 netters retrieving stunned fish, and 1 person with a bucket receiving and disposing of fish. Electrofishing batteries would be recharged using small gasoline powered generators. Based on their previous monitoring efforts, INSE estimated that in a 40 hour work week, 9 sites that were each 100 m long could be completed by a 5 to 6 person crew using the four pass methodology (C. Williams, Institute for Natural Systems Engineering, personal communication with M. Golden, Dixie National Forest, 3/12/2010). Based on this INSE estimate, for UDWR fish removal activities under the Non-chemical Treatment alternative, one removal effort on the 11.5 km mainstem stream (12.8 reaches, 900 m long) on NFS land would require approximately 512 hours (12.8 reaches times 40 hours) or 63 days (8 hours per day) to be completed by a 5 to 6 person crew using the four pass method. An additional effort of approximately 13 days would be needed to treat the 2.3 km mainstem on private property.

Because UDWR's removal activities would need to occur between late-June or early July and September to minimize access, weather, and high stream flow issues, each removal effort would be limited to approximately 20 days to be able to conduct 4 removal efforts in a single year. To be able to treat the entire mainstem stream, on NFS lands and private lands, during any one removal effort, 20 people (four 5-person crews) would be needed. For four removal efforts, this would total up to 80 days per year. As described below, UDWR may need up to 10 years of removal effort under this method.

During the UDWR's 2009 chemical treatment of East Fork Boulder Creek above King's Pasture Reservoir, 23 relatively large inflows or tributaries with permanent fish habitat were identified, along with many smaller springs and seeps capable of at least temporarily holding small fish. These tributaries and inflows varied in length from 10 m to over 750 meters. Additional inflows and tributaries that contain fish habitat are probably present in the reach below Kings Pasture and could add another 30 days or more to the estimated treatment time.

Efficiency of fish removal by electrofishing is substantially lower in certain types of habitats found in the treatment area, especially those with heavy aquatic vegetation, root wads, woody debris, and boulder fields. The time for one removal effort in these types of areas could be higher, and effectiveness could be lower. Also, in order to eliminate the possibility of fish moving between treated and untreated reaches, crews would need to operate simultaneously, which may negatively impact fish-removal efficiency, as stream bed disturbance from upstream crews would impact water clarity and visibility for downstream crews. Because of reduced removal efficiency with electrofishing as the fish removal method, the UDWR project may extend to 10 years.

Post-Fish Removal activities. Post-fish-removal activities by UDWR would be the same as those described for the Proposed Action.

Design Criteria. The following design criteria would be included in the written authorization for use of motorized vehicles off of designated routes and the special use authorization for the burial of removed fish:

- 1. State of Utah decontamination protocols for prevention of the spread of Aquatic Nuisance Species will be followed for all gear and personnel involved with the removal project.
- 2. The Forest Archaeologist will be consulted about potential locations to bury fish to avoid impacts to cultural resources.
- 3. Dead fish collected will be buried no closer than 300 feet from the stream and away from known camping areas to minimize bear/human interactions.
- 4. Access by motorized vehicles will be on National Forest System roads designated for motorized vehicle use to the extent possible. Any use of motorized vehicles off of designated routes will be minimal, and will require written Forest Service approval.
- 5. Trails will be used whenever possible to move from one location to another to minimize soil and vegetation disturbance and to prevent establishing new trails.
- 6. Sensitive plant habitat will be avoided during action implementation.
- 7. Personnel will ensure reach being treated is void of livestock and people not involved with the operation. Treated waters will remain open to fishing.

Actions connected to fish removal actions on NFS lands. The following parts of the UDWR project, as discussed above, are not subject to Forest Service permit requirements, and therefore are not included in the Forest Service decision. They are considered connected actions to UDWR's fish removal activities on NFS lands and thus included in the environmental analysis:

1. As described for the Proposed Action, the UDWR treatment area includes private property, including that owned by Garkane Energy; thus, this area is not under Forest Service jurisdiction.

The expectation is that under the Non-Chemical Treatment alternative, the UDWR would implement non-chemical treatment methods on non-NFS lands, as described below, although the UDWR may decide to use another method or methods to achieve the treatment objective on the private lands or not pursue treatment on the private lands. The flowing portions of the project area on private lands would undergo similar electrofishing removal by UDWR, as described for NFS lands above.

For the non-flowing portions of the project area on private lands, electrofishing would not be effective in removing brook trout from King's Pasture Reservoir or the pond in Kings Pasture. To remove brook trout from these areas without use of chemicals, UDWR would deploy experimental gill nets with many different mesh sizes at several locations and depths throughout each water body. Other studies where this method has been successful at eradicating brook trout suggest that it would take at least two and up to four seasons of semi-continuous netting to eliminate all size classes of trout from small lakes with relatively low trout densities (Knapp and Matthews 1998, Parker et al. 2001).

 Potential recolonization from East Fork Boulder Creek would severely reduce the efficacy of removing brook trout from King's Pasture Reservoir; therefore, UDWR would need to construct a fish migration barrier in East Fork Boulder Creek on private property above King's Pasture Reservoir.

The barrier would generally consist of a small check dam constructed of boulders and large rocks, creating a vertical drop of approximately 5 ft on the downstream side. The location for the barrier would be selected by UDWR to utilize any naturally occurring drops which can be enhanced and where the stream channel and floodplain are confined to minimize the size of the structure and the amount of water impounded behind it. Barrier construction would comply with laws, regulations, and permitting requirements of the State Engineer for stream channel alteration. Barrier materials would be taken from the ground surface, near the stream. The collection of these materials would not require excavation, stream alteration, or vegetation disturbance. If sufficient material is not available on site, additional materials would be hauled to the barrier site from an approved source.

The barrier location would be selected by UDWR to minimize changes in stream gradient, hydraulic function, and water pooling. In addition, the barrier would be constructed by UDWR adjacent to existing roads where equipment access is acceptable, thus requiring little disturbance to surrounding areas. Riparian vegetation would be disturbed as little as possible during the construction of the barrier, while areas where surface disturbance would occur would be restored to pre-project conditions. The barrier would not be placed in areas of cultural or historic significance or in areas where sensitive, threatened or endangered plants occur. It would be designed to operate under the natural fluctuations of a stream flow without routine maintenance. The barrier would be designed to pose little, if any, threat to the natural stream system or its associated riparian area so that if it were to fail, no damage would result to the stream environment. UDWR's maintenance could include

the adjustment or replacement of individual rock materials, but such work would be minor. The barrier could be removed but only after treatment is determined to be fully successful.

Neither netting nor electrofishing are options for UDWR for removing any non-native trout that may be using the upper portion of the penstock inflow or the lower portion of the pipeline from the West Fork Reservoir during treatment efforts. Shutting off water to these areas until they were completely dry would be the only way to ensure complete eradication; however, this is not feasible (M. Avant, Garkane Energy, personal communication with M. Golden, Dixie National Forest, 4/1/2010). Because of this, the effectiveness of the rest of the treatment would be reduced, contributing to the likelihood of the longer period of treatment.

- 3. Stocking of fish by UDWR would be as described for the Proposed Action.
- 4. As described for the Proposed Action, fishing regulations, including whether or not treated waters would remain open to fishing, is under the jurisdiction of UDWR. The expectation is as described for the Proposed Action.

Literature Cited

- Buktenica, M. W., B. D. Mahoney, S. F. Girdner, and G. L. Larson. 2000. Response of a resident bull trout population to nine years of brook trout removal, Crater Lake National Park, Oregon. Pages 127-132 in D. Schill, S. Moore, P. Byorth, and B. Hamre, editors. *Wild Trout VII: Management in the New Millennium, Are We Ready?* Yellowstone National Park, WY. 284 pp.
- Chamberlain, C.B., and D.K. Hepworth. 2002a. Pine Creek rotenone treatment, 2002. A native cutthroat trout restoration project. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 9 pp. + attachments.
- Chamberlain, C.B., and D.K. Hepworth. 2002b. The treatment of Pine Creek, Fremont River drainage, and Pine Creek Reservoir, 2002. A native cutthroat trout and sport fish enhancement project. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 9 pp. + attachments.
- Chamberlain, C.B., and D.K. Hepworth. 2002c Twitchell Creek and Round Willow Bottoms, 2002: A sport fish and native cutthroat trout restoration project. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 8 pp. + attachments.
- Earle, J.E., J.D. Stelfox, and B. Meagher. 2007. Quirk Creek brook trout suppression project 2004-2006. Alberta Sustainable Resource Development, Fish and Wildlife Division, Calgary, Alberta, Canada. 36pp.
- Finlayson, B. J., R. A. Schnick, R. L. Cailteaux, L. DeMong, W. D. Horton, W. McClay, C. W. Thompson, and G. J. Tichacek. 2000. Rotenone use in fisheries management; administrative and technical quidelines manual.
- Fridell, R.A., M.K. Morvilius, M.A. Schijf, and K.K. Wheeler. 2004. Virgin River Basin 2003 treatment projects. Utah Division of Wildlife Resources, Salt Lake City, UT. Publication No. 04-03. 33 pp.

- Fridell, R.A., M.K. Morvilius, and C.B. Rognan. 2005. Virgin River Basin 2004 treatment projects. Publication No. 05-05. Utah Division of Wildlife Resources Publication Number 05-05. 38 pp.
- Fridell, R.A., and A.H. Rehm. 2006. Virgin River Basin 2005 Treatment Projects Utah Division of Wildlife Resources Publication Number 06-05. 30 pp.
- Hardy, T. B., N. Bouwes, C. Williams, and C. Thomas. 2009a. *Trout Population Monitoring in Boulder Creek: 2008 Results*. Utah Water Research Laboratory, Utah State University, Logan, UT. 11 pp.
- Hardy, T. B., C. S. Williams, C. W. Thomas. 2009b. *Trout Population Monitoring in Boulder Creek: 2009 Results*. Utah Water Research Laboratory, Utah State University, Logan, UT. 13pp.
- Hepworth, D.K., C.B. Chamberlain, and J.E. Whelan. 2001a. Pine Creek rotenone treatment, 2001: A native cutthroat trout restoration project. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 4 pp. + attachments.
- Hepworth, D.K., C.B. Chamberlain, and J.E. Whelan. 2001b. Twitchell Creek, Long Willow Bottoms and Round Willow Bottoms, 2001: A native cutthroat trout restoration project. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 4 pp. + attachments.
- Hepworth, D.K., C.B. Chamberlain, and J.E. Whelan. 2001c. West Fork Boulder Creek rotenone treatment, 2001: A native cutthroat trout restoration project. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 9 pp. + attachments.
- Knapp, R. A., and K. R. Matthews. 1998. Eradication of nonnative fish by gill netting from a small mountain lake in California. Restoration Ecology 6: 207-213.
- Kulp, M. A., and S. A. Moore. 2000. Multiple electrofishing removals for elimination rainbow trout in a small southern Appalachian stream. North American Journal of Fisheries Management 20: 259-266.
- Meronek, T. G., P. M. Bouchard, E. R. Bukner, T. M. Burri, K. K. Demmerly, D. C. Hatleli, R. A. Klumb, S. H. Schmidt, and D. W. Coble. 1996. A review of fish control projects. North American Journal of Fisheries Management 16:63-74.
- Meyer, K. A., J. A. Lamansky, Jr., and D. J. Schill. 2006. Evaluation of an unsuccessful brook trout electrofishing removal project in a small Rocky Mountain stream. North American Journal of Fisheries Management 26:849-860.
- Moore, S. E., M. A. Kulp, J. Hammonds, and B. Rosenlund. 2005. *Restoration of Sam's Creek and an Assessment of Brook Trout Restoration Methods, Great Smoky Mountains National Park*. Technical Report/NPA/NRWRD/NRTR-2005/342 U.S. Department of the Interior National Park Service, Water Resources Division, Fort Collins, CO. 36 pp.
- Moore, S. E., G. L. Larson, and B. Ridley. 1986. Population control of exotic rainbow trout in streams of a natural area park. Environmental Management 10: 215-219.
- Ottenbacher, M.J., and D.K. Hepworth. 2001. White Creek rotenone treatment project, August 2001. Utah Division of Wildlife Resources, Southern Region, Cedar City, UT. 6 pp. + attachments.

- Parker, B. R., D. W. Schindler, D. B. Donald, and R. S. Anderson. 2001. The effects of stocking and removal of a nonnative salmonid on the plankton of an alpine lake. Ecosystems 4: 334-345.
- Peterson, D. P., K. D. Fausch, and G. C. White. 2004. Population ecology of an invasion: effects of brook trout on native cutthroat trout. Ecological Applications 14:754-772.
- Shepard, B. B., R. Spoon, and L. Nelson. 2002. A native westslope cutthroat population responds positively after brook trout removal. Intermountain Journal of Science 8:191-211.
- Sousa, R.J., F.P. Meyer, and R.A. Schnick. 1987. Better Fishing through Management. U. S. Fish and Wildlife Service, Washington, D. C.
- Thompson, P. D. and F. J. Rahel. 1996. Evaluation of depletion-removal electrofishing of brook trout in small Rocky Mountain streams. North American Journal of Fisheries Management 16: 332-339.
- US Environmental Protection Agency (EPA). 2007. Registration Eligibility Decision for Rotenone. EPA 738-R-07-005. 44 pp.
- US Environmental Protection Agency (EPA). 1999. Alternative Disinfectants and Oxidants Guidance Manual. EPA 815-R-99-014. 346 pp.

Figure 1. Project area location

